



Evaluation of defoliation on leaf water relations, chlorophyll content, and grain yield of triticale (*x triticosecale* wittmack) genotypes under water stress

E. Bijanzadeh¹, Y. Emam^{2*}

¹College of Agriculture and Natural Resources of Darab, Shiraz University, I. R. Iran

²Department of Crop Production and Plant Breeding, College of Agriculture, Shiraz University, Shiraz, I. R. Iran

* Corresponding Author: yaemam@shirazu.ac.ir

ARTICLE INFO

Article history:

Received 8 June 2015

Accepted 6 June 2016

Available online 5 December 2016

Keywords:

Chlorophyll content

Initial water content

Rate of water loss

Source size

ABSTRACT—Optimizing the source size and its utilization by the sink is one of the main factors enhancing the yield potential and decreasing water demand in crops when exposed to drought. To investigate the effect of defoliation on leaf water relations, chlorophyll content and yield components of five triticale genotypes including Sanabad, Juanillo, ET-83-3, ET-84-5 and ET-84-8 under well-watered (100% FC) and water stress (50% FC) conditions, a controlled experiment was carried out at Shiraz University in 2013. The results showed that ET-84-8 and Sanabad genotypes had higher chlorophyll content (ranged from 49.1 to 54.6 SPAD unit) under water stress. Among the triticale cultivars, water stress caused 21 to 42% decline in the rate of water loss (RWL). In all genotypes except ET-83-3 and Juanillo, the excised leaf water retention (ELWR) slowly decreased under water stress conditions. In all triticale genotypes except ET-84-8, water stress declined main shoot yield 21-22%, while in ET-84-8, it was only 9%. Interestingly, in ET-84-8, grain number per spike was not affected by moisture regimes. Sanabad cultivar, with 2.57 g/g had the highest initial water content (IWC) at defoliation of all leaves except the flag leaf and penultimate leaf treatment under water stress. Under defoliation and water stress, ET-84-8 and Sanabad genotypes showed a greater 100-grain weight ranged from 3.60 to 3.74 g. It was concluded that triticale cultivars were more sink-limited especially under water stress, and source restriction by defoliation which had less effects on main shoot yield could be used as a useful tool for lowering water consumption during grain filling.

INTRODUCTION

Triticale (*X Triticosecale* Wittmack) is an amphiploid hybrid between the female parent wheat (*Triticum aestivum* L.) and the male parent rye (*Secale cereals*) (Ammar et al., 2004). Pfeiffer (2003) suggested that under drought stress conditions and problematic soil regions, triticale showed distinct yield superiority and had adaptive advantages over wheat. Giunta et al. (1993) evaluated durum wheat and triticale genotypes under different moisture regimes in a typical Mediterranean climatic region and observed that grain yield of durum wheat reduced significantly under drought stress, while triticale had a slight and non-significant reduction in grain yield compared to the irrigation control. Triticale was found to have superior tolerance to low nutrient availability, water stress, frost, soil acidity, aluminium and other elemental toxicities and salinity (Lelley, 2006). The objective in the synthesis of triticale was to combine the desirable characteristics of the two species (Oettler, 2005).

Photosynthetically active areas of plant (source) and storage capacity of the grains (sink) after flowering are

the main factors limiting grain yield of cereals (Bijanzadeh and Emam, 2011). Production and partitioning of dry matter in cereals is highly related to sink-source relationships under various environments (Bijanzadeh and Emam, 2010; Zhenlin et al., 1998). The control of grain filling has often been considered in terms of the supply of photosynthate (source limitation) or the capacity of the grain to accumulate available carbohydrate (sink limitation). Carbohydrates for grain filling are supplied concurrently from photosynthetic activity and from temporary storage reserves (Bijanzadeh and Emam, 2011). Sink capacity is a function of the number of grains per unit land area and their potential size (Schnyder, 1993).

Many studies have been reported in which the supply of assimilate per grain is modified by defoliation, shading, thinning or ear manipulation treatments to investigate whether grain filling is sink or source limited (Bijanzadeh and Emam, 2010; Dreccer et al., 1997). Ahmadi et al. (2009) reported that defoliation might change the photosynthetic characteristics of remaining leaves. Also, Joudi et al. (2006) showed that defoliation of wheat increased chlorophyll content and

net photosynthetic rate of remained leaves, however, the rate of increasing depended on the type of cultivars.

In Southern parts of Iran, limited rainfall and water stress occur frequently after anthesis, and water stress is an important limiting factor which can cause major loss in crop productivity. Optimizing the source size after anthesis and its utilization by the sink is one of the major factors enhancing the yield potential in crops especially under water deficit conditions (Bijan-zadeh and Emam, 2011). Ahmadi and Joudi (2007) reported that wheat grain yield is reduced depending on the degree of water deficit and defoliation intensity. Effect of source restriction on yield response and leaf water relation of triticale cultivars has not yet been fully understood, especially under drought stress in Iran. To decrease water consumption during grain filling of triticale by defoliation, this study was undertaken to investigate the effect of source manipulation on leaf water relations, chlorophyll content and yield components of triticale cultivars under water stress.

MATERIALS AND METHODS

A pot experiment was carried out to investigate the effect of defoliation intensity and water stress on leaf water relations, chlorophyll content and yield components of triticale cultivars at the greenhouse of College of Agriculture and Natural Resources of Darab, Shiraz University, Shiraz, Iran, during 2013. Experimental treatments were triticale genotypes, moisture regimes and defoliation. The experimental design was a completely randomized (RCD) one with four replicates. Five triticale genotypes including Sanabad, Juanillo, ET-83-3 (VICUNA_4/ERIZO7//YOGUI_1/GIRAF3/FARAS), ET-84-5 (ERIZO_15/FAHAD_3/POLLMER_2.1) and ET-84-8(ERIZO_6/NIMIR_4/VICUNA_4/3MANATI_1) were sown in 10 kg pots filled with a silty loam soil (8% sand, 68.3% silt, and 23.7% clay, bulk density=1.3 g/cm³, pH= 7.6, %OC=0.051, %N=0.16, P=35 mg kg⁻¹, K=590 mg kg⁻¹ and EC=0.44 dS m⁻¹) and 22 mg kg⁻¹ nitrogen was applied as urea fertilizer. For each cultivar, ten uniform seeds were sown in each 5kg pot (25cm width ×35cm height), and thinned to five seedlings at two-leaf stage. The greenhouse temperature was 25±5°C, with 70±10% relative humidity, and light intensity varied in the range of 650-750 μmol m⁻² s⁻¹.

Before planting, the field capacity (FC) of the soil was determined (FC=28%) in the laboratory to set the moisture regimes as well-watered (100% FC) and water stress (50% FC). Pots were weighted every other day and irrigated according to 100% FC for well-watered treatment and 50% FC for water stress, from booting stage (ZGS45) to physiological ripening.

The source manipulation treatments included defoliation of all leaves (D₁), defoliation of all leaves except the flag leaf (D₂), and defoliation of all leaves except the flag leaf and penultimate leaf (D₃) and control (C). Plants were defoliated at booting stage (ZGS45; Zadoks et al., 1974). To avoid contribution of tillers in grain filling of main shoot, all plants were detilled (Savin and Slafer, 1994).

To determine the RWC, at 10 days after anthesis (10DAA), after cutting the flag leaf of each plant and to avoid water loss, it was sealed in a plastic bag and transferred to the laboratory to measure fresh weight within 1 h after excision. Turgid weight was obtained after soaking leaves in distilled water for 5 h at room temperature (25±2°C). After soaking, leaves were blotted dry with tissue paper to determine turgid weight. Also, dry weight of leaf samples was obtained after oven drying for 72°C. Finally, the RWC was calculated according to Beadle et al.'s (1993) method by:

$$RWC = \frac{(W_0 - W_d)}{(W_t - W_d)} \times 100$$

Initial water content (IWC) was calculated as:

$$IWC = \frac{(W_0 - W_d)}{(W_d)}$$

w₀= fresh weight (g), W_d= dry weight (g) of leaves placed in an oven at 50° C for 24 h and re-weighed. W_t=turgid weight(g) (Yang et al., 1991).

To measure the rate of water loss (RWL), the flag leaves were collected and weighed (W1). Then, the leaves were wilted at 30°C for 24 h and re-weighed (W2) and transferred to an oven at 50° C for 24 h and then weighed (W3). RWL was calculated by the equation suggested by Yang et al. (1991):

$$RWL = \left(\frac{W1 - W2}{W3} \right) \left(\frac{t1 - t2}{60} \right)$$

Where t₁ and t₂ are the time of measurement for initial and wilted weight (in minutes).

Also, for determining excised leaf water retention (ELWR), the flag leaves were collected and weighed, then kept at 30°C for 5 hours and reweighed. ELWR was then calculated using the following equation (Wang and Clarke, 1992):

The SPAD meter was used to estimate chlorophyll content non-destructively (Barraclough and Kyte, 2001). It was a hand-held spectrometer which measured light (650 nm) absorbed by a single leaf. In this study, the SPAD chlorophyll meter (Opti-Sciences X. USA) was used to acquire a rapid estimate of flag leaf chlorophyll content.

$$ELWR = \left(1 - \frac{\text{weight of fresh leaves} - \text{weight of leaves after 5 hours}}{\text{weight of fresh leaves}} \right) \times 100$$

Water-related variables and the chlorophyll content were measured at anthesis stage (ZGS 64). The water related parameters were measured on three plants randomly selected from each pot.

Finally, plants were harvested at physiological maturity, oven-dried at 72 °C, and then main shoot grain yield and yield components including grain number per spike and 100-grain weight(g) were determined. The collected data were subjected to the analysis of variance by SAS 9.1 software (2002) and the means were separated using Fisher's LSD protected test at 0.05 probability level.

RESULTS AND DISCUSSION

Chlorophyll Content

Interaction effects of genotype \times moisture regimes \times defoliation on Chlorophyll content of flag leaf was significant at 0.05 probability level (data not shown). ET-84-8 and Sanabad genotypes showed higher chlorophyll content (ranged from 49.1 to 54.6 SPAD unit) compared to other cultivars when exposed to water stress (Fig.1). In contrast, water stress was associated with a decline in chlorophyll content of ET-84-5 and ET-83-3 cultivars. Ahmadi and Joudi (2007) reported similar changes in chlorophyll content among wheat cultivars. Furthermore, in a greenhouse experiment, Barraclough and Kyte (2001) reported that chlorophyll content of remained leaf of winter wheat (CV. Hereward) decreased significantly under water stress. In our study, it appeared that accelerated senescence of flag leaf and penultimate leaf under water stressed due to the decrease of water content, and reduced chlorophyll content of flag leaf and penultimate leaf, compared to well-watered conditions.

Leaf Water Characteristics

In all triticale genotypes at 10 days after anthesis (DAA), water stress caused a significant reduction in relative water content of flag leaf (RWC) at the similar level of

source restriction treatments (Table 1). In D₃ treatment, RWC in ET-83-3 decreased sharply from 71.4 to 38.4 (46.2% reduction) more than the other genotypes under water stress. Also, in all genotypes, no significant differences were found between D₂ and D₃ treatments and the presence of the penultimate leaf had no effect on RWC of flag leaf.

Bijan-zadeh and Emam (2010) reported that some wheat cultivars with lower transpiration rate might maintain higher RWC in their leaves under water stress conditions. They also reported that source restriction by removal of transpiring leaves which were less effective in grain filling could be a useful tool to decrease water loss during grain filling under dry and semi-dry area conditions. Ahmadi and Joudi (2007) found that RWC of flag leaf was not affected by source restriction and defoliation increased transpiration rate of remaining flag leaf and bare soil evaporation as soil was exposed to sunlight. However, Matin (1999) reported that drought tolerant cultivars of barley usually maintained higher leaf RWC under drought stress. Similar to our results, Lonbani and Arzani (2011), too, reported highly significant differences among genotypes for RWC so that 'Moreno' and 'Prego' triticale cultivars showed the highest and Zoro' cultivars had the lowest RWC under water stress conditions. Schonfeld et al. (1998) observed a decline in the amount of RWC in wheat due to water stress and reported a higher RWC for the tolerant genotypes. Accordingly, ET-84-8 and Sanabad triticale genotypes were ranked as drought tolerant while Juanillo and ET-83-3 ranked as drought sensitive genotypes (Table 1).

For each moisture regime, the amount of initial water content (IWC) was not affected by defoliation treatment and water stress decreased IWC in Juanillo and ET-83-3 triticale genotypes more than Sanabad and ET-84-8 (Table 1). On the other hand, D₂ and D₃ treatments of Sanabad with 2.57 and 2.41 g/g had higher IWC under water stress, respectively.




Fig. 1. Effect of moisture regimes and defoliation on flag leaf chlorophyll content (SPAD unit) of triticale cultivars at 10 days after anthesis (DAA). Control(c), defoliation of all leaves except the flag leaf (D2), and defoliation of all leaves except the flag leaf and penultimate leaf (D3). Vertical bars represent SE.

Table 1. Effect of moisture regimes and defoliation on relative water content (RWC), initial watercontent (IWC), rate of water loss (RWL), and excised leaf water retention (ELWR) of flag leaf of triticale cultivars at 10 days after anthesis (DAA).

Barley cultivars	Moisture regime	Source restriction	RWC (%)	IWC (g/g)	RWL (g/g.h)	ELWR (%)
Sanabad	Well-watered	C [†]	76.8	2.61	0.53	83
		D2	77.3	2.78	0.57	80
		D3	77.1	2.63	0.61	81
	Water stress	C	60.2	2.43	0.38	80
		D2	61.1	2.57	0.31	79
		D3	62.4	2.41	0.37	77
Juanillo	Well-watered	C	74.3	1.98	0.67	76
		D2	75.6	1.93	0.72	79
		D3	75.5	1.82	0.73	75
	Water stress	C	48.7	1.14	0.56	61
		D2	50.3	1.21	0.50	62
		D3	48.1	1.11	0.51	60
ET-83-3	Well-watered	C	71.2	1.73	0.76	83
		D2	72.3	1.77	0.84	80
		D3	71.4	1.56	0.83	86
	Water stress	C	41.3	1.09	0.61	59
		D2	39.2	1.13	0.56	55
		D3	38.4	1.11	0.53	57
ET-84-5	Well-watered	C	71.3	2.03	0.78	85
		D2	72.4	2.01	0.84	86
		D3	72.0	1.98	0.86	81
	Water stress	C	40.4	1.17	0.67	78
		D2	41.3	1.26	0.59	77
		D3	43.2	1.20	0.61	73
ET-84-8	Well-watered	C	78.3	1.96	0.51	86
		D2	80.1	2.11	0.58	84
		D3	79.2	1.94	0.50	83
	Water stress	C	63.2	1.88	0.33	81
		D2	65.2	1.90	0.30	83
		D3	64.1	1.84	0.36	82
LSD (0.05)			5.7	0.33	0.16	9

[†]Control (c), defoliation of all leaves except the flag leaf (D2), and defoliation of all leaves except the flag leaf and penultimate leaf (D3)

Among the triticale cultivars, water stress caused 21 to 42% decline in the rate of water loss (RWL). This may indicate some inhibiting mechanisms of water loss under water stress in all triticale genotypes (Table 1). Overall, under water stress, ET-84-8 and Sanabad decreased significantly RWL more than the other genotypes. GolestaniAraghi and Assad (1998) reported a significant decline in the RWL of wheat under water stress conditions. In a similar study, Lonbani and Arzani (2011) reported that triticale genotypes had lower RWL than wheat genotypes. This may indicate more efficient use of water by triticale genotypes compared to wheat genotypes. Wang and Clark (1992) indicated that parameters measured in the excised leaves at minimum stomatal aperture, such as IWC, RWL, and epidermal conductance showed genotypic differences in wheat and this might be related to yield in dry land environments. In contrast to our results, Haley et al. (2002) reported that the amounts of RWL for dry land and irrigated wheat were not different. Furthermore, Jaradat and

Konzak (2003) suggested greater RWL for wheat genotypes in stressed environments.

In all genotypes except ET-83-3 and Juanillo, the excised leaf water retention (ELWR) decreased slowly when exposed to water stress (Table 1). Also, ELWR of ET-84-8, Sanabad and ET-84-5 ranged from 73% to 83% and were higher than Jaunillo and ET-84-3 significantly. Lonbani and Arzani (2011) declared that water stress increased excised leaf water retention (ELWR); this phenomenon showed that the probable mechanism for water retention in the leaf under stress conditions might be leaf rolling or the decrease in leaf area. They concluded that among the plant water relation parameters, ELWR could be a superior indirect selection criterion for grain yield. Wenzel (1997) observed that EWLW of sorghum genotypes were different under drought stress conditions, and concluded that EWLW was a satisfactory screening technique for drought resistance in sorghum genotypes.

Grain Yield and Yield Components

Grain number per spike responded significantly to defoliation treatment in Juanillo and ET-83-3 (Table 2).

Interestingly, in ET-84-8 cultivar, grain number per spike was not affected by moisture regimes and D2 treatment with 46.5 grains per spike had the highest grain number per spike, compared to other genotypes, under water stress. In a similar study, Alam et al. (2008) reported that defoliation of all leaves except the flag leaf after anthesis decreased the grain number per spike in wheat cultivars. In all triticale cultivars, water stress declined main shoot yield significantly; however, in ET-84-8 and Sanabad, the decrease rate was less than that of the other cultivars (Table 2). Overall, in each moisture regime, grain yield was not affected by source

restriction treatments. It has previously been demonstrated that grain cereals might be more sink-limited, especially under water stress conditions (Emam and Seghatoleslami, 2005). Joudi et al., (2006) also, reported lower response of grain yield to changes in assimilate availability by source restriction. They concluded that this might be related to more sink limitation in some bread wheat cultivars. Under water stress and defoliation treatments, maximum main shoot yield was observed in ET-84-8 and Sanabad which ranged from 1.62 to 1.73 g. In ET-84-8 and Sanabad cultivars, D₁ treatment reduced main shoot yield only 6.1 and 5.7%, respectively, when plants were exposed to water stress, while main shoot yield in D₁ treatment of Juanillo cultivar reduced 15.6%.

Table 2. Interaction effects of moisture regimes and defoliation on yield components of triticale genotypes under well-watered and water stress (50% FC) conditions.

Barley cultivars	Irrigation regime	Source restriction	Grain number per spike	100-grain weight(g)	Main shoot yield(g)
Sanabad	Well-watered	C	47.1	4.12	1.94
		D1	43.3	3.99	1.72
		D2	46.5	4.01	1.86
		D3	46.6	4.03	1.87
	Water stress	C	45.7	3.77	1.72
		D1	45.1	3.60	1.62
		D2	45.4	3.72	1.68
		D3	45.3	3.74	1.69
Juanillo	Well-watered	C	44.6	4.1	1.82
		D1	40.1	3.88	1.55
		D2	40.3	3.96	1.59
		D3	40.8	3.99	1.62
	Water stress	C	40.4	3.50	1.41
		D1	36.1	3.31	1.19
		D2	37.8	3.38	1.27
		D3	37.9	3.39	1.28
ET-83-3	Well-watered	C	42.1	4.01	1.68
		D1	37.1	3.81	1.41
		D2	38.2	3.93	1.50
		D3	38.9	3.98	1.54
	Water stress	C	37.1	3.41	1.26
		D1	34.2	3.21	1.09
		D2	35.1	3.28	1.15
		D3	35.9	3.30	1.18
ET-84-5	Well-watered	C	40.4	4.04	1.63
		D1	38.1	3.78	1.44
		D2	38.3	3.84	1.47
		D3	38.4	3.86	1.48
	Water stress	C	36.2	3.33	1.20
		D1	32.8	3.11	1.02
		D2	35.3	3.23	1.14
		D3	35.6	3.29	1.19
ET-84-8	Well-watered	C	48.2	4.14	1.99
		D1	45.6	4.01	1.86
		D2	46.1	4.09	1.88
		D3	46.8	4.11	1.92
	Water stress	C	47.1	3.78	1.78
		D1	46.3	3.62	1.67
		D2	46.5	3.71	1.72
		D3	46.4	3.73	1.73
LSD (0.05)			1.2	0.11	0.13

^aControl (c), defoliation of all leaves (D1), defoliation of all leaves except the flag leaf (D2), and defoliation of all leaves except the flag leaf and penultimate leaf (D3).

Our finding regarding Et-84-8 and Sanabad cultivars was in agreement with that of Emam and Dastfal (1997) who showed that cultivars with higher grain yield potential under well-watered condition had also higher grain yield under water stress as well. Interestingly, in all cultivars, penultimate leaf had no significant effect on main shoot yield (comparison of D₂ with D₃ treatments; Table 2). Guttieri et al. (2001) observed that moisture deficit induced reduction in 100-grain yield of wheat due to reduction in grain weight while the differential effect of moisture deficit on specific cultivars could be due to reduction in the number of grains per spike. In a similar study, Fayaz and Arzani (2011) reported that grain number per spike of triticale cultivars ranged from 2.84 for Roshan to 5.27 for Moreno under well-watered conditions and ranged from 2.27 for Roshan to 3.56 for Lasko under water stress conditions. Also, 100-grain weight varied from 2.89 g to 4.56 g for Prego and Alamos83 cultivars under well-watered conditions, respectively. Under water stress conditions, 100-grain weight decreased significantly and ranged from 2.23 g for Prego to 3.54 g for Alamos83. The lowest and highest reduction in grain yield due to water stress were also observed in Alamos83 (24%) and Prego (65.4%) cultivars, respectively. In spite of our results, Riaz and Choudhry (2003) reported that genotypes with higher 100-grain weight under well-watered conditions may not be superior in this trait under water stress conditions. Indeed, the limitation of moisture might force the plant to complete its grain filling in a relatively shorter duration (Emam and Seghatoleslami, 2005). The relationship between RWC and main shoot yield at 10DAA ($R^2=0.73$) was highly

significant (Fig. 2a). Also, a significant positive relationship was found between flag leaf chlorophyll content and main shoot yield ($R^2=0.84$) (Fig. 2b). In a similar study, Bijanzadeh and Emam (2010a) reported a positive relationship between grain yield with RWC ($R^2=0.78$) and chlorophyll content ($R^2=0.61$) at 10DAA in five wheat cultivars.

CONCLUSIONS

Our results revealed that ET-84-8 and Sanabad genotypes might be suitable for source restriction and water stress conditions, since they did not show any significant reduction in 100-grain weight, number of grains per spike, and grain yield, when the assimilate availability was reduced by defoliation treatment. However, yield and yield components of ET-83-3, ET-84-5 and Juanillo cultivars were sensitive to water stress, indicating that in these cultivars reduced assimilate availability by water stress was associated with a decrease in sink size. Additionally, flag leaf water relations including RWC, IWC, RWL, and ELWR of ET-83-3 and Juanillo cultivars were more sensitive to water deficit, while ET-84-8 and Sanabad maintained higher chlorophyll content, RWC, RWL, and ELWR under water stress. It could be recommended that the selection and growing of cultivars such as ET-84-8 and Sanabad with small responses to source restriction, might be a good approach for yield potential improvement, particularly in regions with restricted water availability later in the growing season.




Fig. 2. Relationship between grain yield of triticale cultivars with RWC (a) and flag leaf chlorophyll content (b).

REFERENCES

- Ahmadi, A., & Joudi, M. (2007). Effects of timing and defoliation intensity on growth, yield, and gas exchange rate of wheat grown under well-watered and drought conditions. *Pakistan Journal of Biological Science*, 10, 379-380.
- Alam, M.S., Rahman, A.H., Nesa, M.N., Khan, S.K., & Siddiquie, N.A. (2008). Effect of source and/or sink restriction on the grain yield in wheat. *Journal of Applied Science Research*, 4, 258-261.
- Ammar, K., M., Mergoum, & S., Rajaram., (2004). The history and evolution of triticale. Pp. 1-9. In M., Mergoum, & H. Gomez-Macpherson (1sted). *Triticale Improvement and Production*. (pp.1-9). Rome, Italy.
- Barracough, P.B., & Kyte, J. (2001). Effect of water stress on chlorophyll meter readings in winter wheat. In W.J., Horst, M.K., Schenk, A., Burkert, N., Claassen&Flessa, H. (Ed.). *Plant Nutrition- Food Security and Sustainability of Agro-Ecosystems*, (pp. 722-766), Kluwer Academic Publishers, Netherlands.
- Beadle, C.L., Ludlow, M.M., & Honeysett, J.L. (1993). *Photosynthesis and production in a changing environment*. Chapman and Hall, London. Pp. 113-128.
- Bijan-zadeh, E., & Emam, Y. (2010). Effect of defoliation and drought stress on yield components and chlorophyll content of wheat. *Pakistan Journal of Biological Science*, 13, 699-705.
- Bijan-zadeh, E., & Emam, Y. (2011). Evaluation of assimilate remobilization and yield of wheat cultivars under different irrigation regimes in arid climate. *Archives of Agronomy and Soil Science*, 58, 1243-1259.
- Borras, L., Slafer, G.A., & Otegui, M.E. (2004). Seed dry weight response to source-sink manipulations in wheat, maize and soybean: A quantitative reappraisal. *Field Crops Research*, 86, 131-146.
- Dreccer, M.F., Grashoff, C., & Rabbinge, R. (1997). Source-sink ratio in barley (*Hordeumvulgare* L.) during grain filling: effect on senescence and grain protein concentration. *Field Crops Research*, 49, 269-277.
- Emam, Y., & Seghatoleslami, M.J. (2005). *Crop yield, physiology and processes*. Shiraz University Press. 593 pp.(In Persian)
- Emam, Y., & Dastfal, M. (1997). Above and below ground responses of winter barley plants to chlormequat in moist and drying soil. *Crop Research*, 14, 457-470.
- Fayaz N., & Arzani A., 2011. Moisture stress tolerance in reproductive growth stages in triticale (X *Triticosecale* Wittmack) cultivars under field conditions *Crop Breeding Journal*, 1, 24-34. (In Persian)
- Giunta, F.R. Motzo, & Deidda, M. (1993). Effect of drought on yield and yield components of durum-wheat and triticale in a Mediterranean environment. *Field Crops Research*, 33, 399-409.
- GolestaniAraghi, S., & Assad, M.T. (1998). Evaluation of four screening techniques for drought resistance and their relationship to yield reduction ratio in wheat. *Euphytica*, 103,293-299.
- Guttieri, M.J., Stark, J.C. Brien, K.O., & E. Souza, (2001). Relative sensitivity of spring wheat grain yield and quality parameters to moisture deficit. *Crop Science*, 41, 327-335.
- Haley, S.D., Quick, J.S., & Morgan, J. A. (2002). Excised-leaf water status evaluation and associations in field-grown winter wheat. *Canadian Journal of Plant Science*, 83, 55-63.
- Jaradat, A., & Konzak, C.F. (2003). Screening of wheat genotypes for drought tolerance. I. Excised-leaf water retention. *Cereals Research Communication*, 31, 179-187.
- Joudi, M., Ahmadi, A., Poustini, K., & Sharifzadeh, F. (2006). Effect of leaf elimination on the effectiveness of flag leaf photosynthesis and seed growth in bread wheat. *Iranian Journal of Agriculture Science*, 2, 203-211.
- Lelley, T. (2006). Triticale: A low-input cereal with untapped potential. In: Singh, R.J., & Jauhar, P. P. (Eds.) *Genetic resources chromosome engineering and crop improvement*. (pp. 395-430). Boca Raton, CRC Press, Taylor & Francis Group, FL.
- Lonbani M., & Arzani A. (2011). Morpho-physiological traits associated with terminal drought stress tolerance in triticale and wheat. *Agriculture Research*, 9, 315-329.
- Matin, M.A., Jarvis, H.B., & Hayden, F. (1999). Leaf water potential, relative water content and diffusive resistant as screening techniques for drought tolerance in barley. *Agronomy Journal*, 91,100-105.
- Oettler, G. (2005). Centenary review. The fortune of a botanical curiosity-triticale: Past, Present and future. *Journal of Agriculture Science*, 143, 329-346.
- Pfeiffer, W.H. (2003). Triticale improvement strategies at CIMMYT: Exploiting adaptive patterns and end-use orientation. *Triticale Topics*, 21, 18-27.
- Riaz, R., & Choudhry, M.A. (2003). Genetic Analysis of some economic traits of wheat under drought condition. *Asian Journal of Plant Science*, 2, 790-796.
- Savin, R., & Slafer, G.A. (1994). Shading effects on the yield of an Argentina wheat cultivar. *Journal of Agriculture Science*, 116,1-7.
- Schnyder, H. (1993). The role of carbohydrate storage and redistribution in the source-sink relations of wheat and barley during grain filling. *New Phytologica*, 123, 233-245.
- Schonfeld, M. A., Johnson, R.C., Carver B.F., & Mornhinweg, D.W. (1998). Water relations in winter wheat as drought resistance indicators. *Crop Science*, 28, 526-531.
- Wang, H., & Clarke, M. (1992). Relationship of excised-leaf water loss and stomatal frequency in wheat. *Canadian Journal of Plant Science*, 73, 93-99.
- Wenzel, W.G. (1997). Leaf water retention of excised leaves as a measure of drought resistance in grain sorghum (*Sorghum bicolor* L.) genotypes. *South African Journal of Plant Science*, 14:31-34.
- Yang, R.C., Jana, S., & Clark J.M. (1991). Phenotypic diversity and associations of some potentially drought-responsive characters in durum wheat. *Crop Science*, 31, 1484-1491.
- Zadoks, J.C., Chang, T.T., & Konzak, C.F. (1974). A decimal code for the growth stages of cereals. *Weed Research*, 14:11-16.
- Zhenlin, W., Yanping, Y., Mingrong, H., & Hongmingand, C. (1998). Source-sink manipulation effect on post anthesis and grain setting on spike in winter wheat. *Photosynthetica*, 35, 453-459.



ارزیابی روابط آبی برگ، محتوای کلروفیل و عملکرد دانه ژنوتیپ های تریتیکاله در شرایط کم آبی و برگ زدایی

احسان بیژن زاده^۱، یحیی امام^{۲*}

^۱ دانشکده کشاورزی و منابع طبیعی داراب، دانشگاه شیراز، داراب، ج.ا.ایران
^۲ گروه زراعت و اصلاح نباتات، دانشکده کشاورزی، دانشگاه شیراز، شیراز، ج.ا.ایران.

*نویسنده مسئول

اطلاعات مقاله

تاریخچه مقاله:

تاریخ دریافت: ۱۳۹۴/۹/۳

تاریخ پذیرش: ۱۳۹۵/۳/۱۷

تاریخ دسترسی: ۱۳۹۵/۹/۱۵

واژه‌های کلیدی:

مقدار کلروفیل

مقدار آب اولیه برگ

سرعت از دست دادن آب برگ

اندازه مبدا

چکیده- بهینه کردن اندازه مبدا و استفاده مواد پروده آن در مقصد یکی از عوامل مهم در افزایش پتانسیل عملکرد و کاهش تقاضای آب در گیاهان زراعی در شرایط تنش خشکی می باشد. بمنظور بررسی اثرات برگ زدایی بر روابط آبی برگ پنج ژنوتیپ تریتیکاله شامل سناباد، جوانیلو، ET-۸۳-۳، ET-۸۴-۵، ET-۸۴-۸ و ET-۸۴-۸ در شرایط آبیاری مطلوب (۱۰۰ درصد ظرفیت مزرعه) و تنش کم آبی (۵۰ درصد ظرفیت مزرعه)، آزمایشی گلخانه‌ای در سال ۱۳۹۲ در دانشگاه شیراز به اجرا در آمد. نتایج نشان داد که رقم ET-۸۴-۸ و سناباد محتوی کلروفیل بالاتری (در دامنه از ۴۹/۱ تا ۵۴/۶ واحد اسپد) در شرایط تنش کم آبی داشتند. در بین ارقام تریتیکاله تنش کم آبی باعث کاهش ۲۱ تا ۴۲ درصدی در سرعت از دست دادن آب برگ (RWL) شد. در همه ارقام به غیر از ET-۸۳-۳ و جوانیلو مقدار آب نگهداری شده در برگ (ELWR) به آرامی در شرایط تنش کم آبی کاهش یافت. در همه ارقام تریتیکاله به غیر از ET-۸۴-۸، تنش کم آبی عملکرد ساقه اصلی را بین ۲۱ تا ۲۲٪ کاهش داد در حالی که این کاهش در رقم ET-۸۴-۸ تنها ۹٪ بود. جالب اینکه تعداد دانه در سنبله در رقم ET-۸۴-۸ تحت تأثیر رژیم رطوبتی قرار نگرفت. همچنین وقتی همه برگ‌ها به غیر از برگ پرچم و برگ ماقبل پرچم حذف شدند، رقم سناباد با ۲/۵۷ گرم بر گرم دارای مقدار آب اولیه برگ (IWC) زیادتری بود. در شرایط تنش کم آبی و برگ زدایی، ارقام ET-۸۴-۸ و سناباد وزن هزار دانه بیشتری (در دامن‌های از ۳/۶ تا ۳/۷۴ گرم) داشتند. می توان نتیجه گرفت که ارقام تریتیکاله بویژه در شرایط کم آبی مقصد- محدود بوده و اعمال محدودیت مبدا از راه برگ زدایی تأثیر کمتری در عملکرد دانه داشت، که این موضوع می‌تواند به عنوان ابزاری سودمند در کاهش آب مصرفی در دوره پر شدن دانه مورد استفاده قرار گیرد.